
Q1 Small mammals and birds help spread Lyme disease because they act as natural carriers for Borrelia.
100.00
80.00
60.00
40.00
20.00
0.00
42.50
90.91
22.50
6.06
35.00
3.03
TRUE
FALSE
Don't know
% Before
% After


Q2 Lyme disease is the most common vector-borne disease in Europe and the USA.
100.00
80.00
60.00
40.00
20.00
0.00
65.00
90.32
12.50
0.00
22.50
9.68
TRUE
FALSE
Don't know
% Before
% After


Q3 Ticks must be attached for at least 24 hrs before the infection can be transmitted?
120.00
100.00
80.00
60.00
40.00
20.00
0.00
7.50
3.23
75.00
96.77
17.50
0.00
TRUE
FALSE
Don't know
% Before
% After


Q4 A child of 9 yrs of age who has a history of tick attachment followed by an erythema migrans rash should be treated with doxycycline.
80.00
60.00
40.00
20.00
0.00
55.00
70.97
20.00
29.03
25.00
0.00
TRUE
FALSE
Don't know
% Before
% After


Q5 Ticks may carry and transmit other pathogens as well as Borrelia.
120.00
100.00
80.00
60.00
40.00
20.00
0.00
85.00
100.00
2.50
0.00
12.50
0.00
TRUE
FALSE
Don't know
% Before
% After


Q6 It takes at least 6 weeks before Borrelia begins to spread around the body.
80.00
60.00
40.00
20.00
0.00
20.00
19.35
45.00
70.97
35.00
9.68
TRUE
FALSE
Don't know
% Before
% After


Q7 An EM rash is diagnostic of lyme disease without the need for a blood test.
120.00
100.00
80.00
60.00
40.00
20.00
0.00
52.50
96.77
37.50
3.23
12.50
0.00
TRUE
FALSE
Don't know
% Before
% After


Q8 Facial palsy is a recognised symptom of Lyme disease.
120.00
100.00
80.00
60.00
40.00
20.00
0.00
57.50
100.00
7.50
0.00
35.00
0.00
TRUE
FALSE
Don't know
% Before
% After


Q9 If the Lyme serology is negative at 12 weeks post bite, then Lyme disease has been excluded.
120.00
100.00
80.00
60.00
40.00
20.00
0.00
12.50
0.00
55.00
100.00
32.50
0.00
TRUE
FALSE
Don't know
% Before
% After


Q10 Women who are treated for Lyme disease during pregnancy can be confidently reassured that there is no risk to their unborn foetus.
100.00
80.00
60.00
40.00
20.00
0.00
0.00
0.00
40.00
80.65
60.00
19.35
TRUE
FALSE
Don't know
% Before
% After